

- 8 Sardana K, Manchanda V, Rajpal M et al. Bacterial pyoderma in children and therapeutic options including management of community acquired methicillin resistant *Staphylococcus aureus*. *Int J Dermatol* 2006 (in press).
- 9 Centers for Disease Control and Prevention. Four pediatric deaths from community acquired methicillin-resistant *Staphylococcus aureus*—Minnesota and North Dakota, 1997–1999. *MMWR Morb Mortal Wkly Rep* 1999; **48**:707–10.
- 10 Fridkin SK, Hageman JC, Morrison M et al. Methicillin-resistant *Staphylococcus aureus* disease in three communities. *N Engl J Med* 2005; **352**:1436–44.

Conflicts of interest: none declared.

- 3 Vandenesch F, Naimi T, Enright MC et al. Community-acquired methicillin-resistant *Staphylococcus aureus* carrying Panton–Valentine leukocidin genes: worldwide emergence. *Emerg Infect Dis* 2003; **9**:978–84.
- 4 Yamaguchi T, Yokota Y, Terajima J et al. Clonal association of *Staphylococcus aureus* causing bullous impetigo and the emergence of new methicillin-resistant clonal groups in Kansai District in Japan. *J Infect Dis* 2002; **185**:1511–16.

Conflicts of interest: none declared.

Community-acquired methicillin-resistant *Staphylococcus aureus*: different populations, different results: reply from authors

DOI: 10.1111/j.1365-2133.2006.07544.x

SIR, We are grateful to Sardana et al. for their interesting comments. They found, in contrast to the current published data and our results,^{1,2} that community-acquired methicillin-resistant *Staphylococcus aureus* (CA-MRSA) was not always responsible for severe skin manifestations. These different clinical presentations may be related to the different *S. aureus* clones involved in skin infections. It is becoming evident that the various CA-MRSA clones circulating in the different parts of the world harbour different antibiotic resistance profiles and toxin genes.³ Therefore according to the clones circulating, the clinical manifestations may differ from one country to another. Most CA-MRSA clones harbour the Panton–Valentine toxin genes which are responsible for pus and necrosis and cause furuncles and abscesses, but other clones harbour exfoliative toxin genes responsible for blisters and outbreaks of impetigo.⁴ Therefore it is important to identify the circulating CA-MRSA clones in a geographical focus, better to characterize the clinical manifestations associated with these clones.

Unité d'Infectiologie–Dermatologie,
Hôpital Bonnet, 83700 Fréjus, France
E-mail: del-giudice-p@frejus-saint-raphael.fr

P.D. DEL GIUDICE

References

- 1 Del Giudice PD, Blanc V, Durupt F et al. Emergence of two populations of methicillin-resistant *Staphylococcus aureus* with distinct epidemiological, clinical and biological features, isolated from patients with community-acquired skin infections. *Br J Dermatol* 2006; **154**:118–24.
- 2 King MD, Humphrey BJ, Wang YF et al. Emergence of community-acquired methicillin-resistant *Staphylococcus aureus* USA 300 clone as the predominant cause of skin and soft-tissue infections. *Ann Intern Med* 2006; **144**:309–17.

A surgical treatment for anxiety-triggered palmar hyperhidrosis is not unlike treating tearfulness in major depression by severing the nerves to the lacrimal glands

DOI: 10.1111/j.1365-2133.2006.07547.x

SIR, The recent study of Krogstad et al.¹ in this Journal, reporting on the daily pattern of sweating and response to stress in patients with palmar hyperhidrosis, is an important contribution to the clinical literature. Patients and controls did not differ while doing 'physical activity' or 'other activity'. In contrast, in patients, but not in controls, palmar hyperhidrosis was most severe at mid-day during the working day (when psychological stress levels are likely to be highest). These authors conclude that their 'finding supports the idea that thermoregulatory sweating is normal in hyperhidrotic patients' but found extensive evidence that psychological stress influences patients with palmar hyperhidrosis more than controls.¹

The surgical literature (see Krogstad et al. for references) advocates sympathectomy or other surgical procedures as the first line of treatment for palmar hyperhidrosis. This is despite the frequent postsympathectomy complications, most commonly truncal compensatory hyperhidrosis, a condition which we contend is at least as uncomfortable and as socially embarrassing as palmar hyperhidrosis. Long-lasting postsurgical compensatory truncal hyperhidrosis has been reported to occur in 44–86% of patients.^{2,3}

Palmar hyperhidrosis of clinical severity is a hallmark physical sign of many anxiety disorders, including generalized anxiety disorder, panic disorder, posttraumatic stress disorder, and especially social phobia.⁴ These are increasingly well understood and highly treatable neurobiological conditions. They are moderately heritable hard-wired fear responses,⁵ and are linked to amygdalar and locus coeruleus hyper-reactivity during psychosocial stress.^{6,7} Anxiety disorders are known to be much more common among women. This is consistent with the finding of Krogstad et al. that among controls sweating was reported more often by men, while among the hyperhidrosis group sweating was reported more often among women.

It may be relevant that a disproportionate percentage of the published surgical studies of palmar hyperhidrosis has been

conducted in Asian countries where a strong stigma exists against having any mental disorder. Because of this, many individuals from these cultures (e.g. Japan, Korea, Taiwan and Hawaii) would rather go under the knife than take psychotropic medications. However, all patients with palmar hyperhidrosis should be offered and strongly encouraged to complete a full course of a psychopharmacological treatment (e.g. with a serotonin/noradrenaline reuptake inhibitor) for anxiety, the underlying cause of palmar hyperhidrosis,^{4,8} before undergoing a surgical procedure. A surgical treatment for anxiety-triggered palmar hyperhidrosis is not unlike treating tearfulness in major depression by severing the nerves to the lacrimal glands. We have recently made a similar argument advocating a psychopharmacological, rather than a surgical, first-line treatment for blushing.⁹

Finally, it cannot be more strongly emphasized that, like major depression, the above anxiety disorders are a risk factor for suicide.⁴ Referring an anxious patient with palmar hyperhidrosis to surgery without first completing a proper trial of psychotropic medication may constitute malpractice especially if the patient experiences some of the more severe surgical complications which can occur during sympathectomy.

Acknowledgments

This material is based upon work supported in part by the Office of Research and Development, Medical Research Service, Department of Veterans Affairs, Pacific Islands Health Care System, Spark M. Matsunaga Medical Center. Support was also provided by a National Alliance for Research on Schizophrenia and Depression (NARSAD) Independent Investigator Award, and the VA National Center for Post-traumatic Stress Disorder.

*National Center for Post-traumatic Stress Disorder, Pacific Islands Division,
Department of Veterans Affairs,
Pacific Islands Health Care System,
Spark M. Matsunaga Medical Center,
1132 Bishop Street #307,
Honolulu 96813 2830, HI, U.S.A.
†Asia-Pacific Center for Biosecurity,
Disaster & Conflict Research, Asia-Pacific
Institute for Tropical Medicine and
Infectious Disease, Department of Tropical Medicine, Medical Microbiology and
Pharmacology, John A. Burns School of Medicine,
University of Hawaii, Honolulu, HI, U.S.A.
E-mail: h.bracha@med.va.gov

References

- 1 Krogstad AL, Mork C, Piechnik SK. Daily pattern of sweating and response to stress and exercise in patients with palmar hyperhidrosis. *Br J Dermatol* 2006; **154**:1118–22.
- 2 Schick CH, Horbach T. Sequelae of endoscopic sympathetic block. *Clin Auton Res* 2003; **13** (Suppl. 1):I36–9.

- 3 Furlan AD, Mailis A, Papagapiou M. Are we paying a high price for surgical sympathectomy? A systematic literature review of late complications. *J Pain* 2000; **1**:245–57.
- 4 den Boer JA. Social anxiety disorder/social phobia: epidemiology, diagnosis, neurobiology, and treatment. *Compr Psychiatry* 2000; **41**:405–15.
- 5 Bracha HS. Human brain evolution and the 'neuroevolutionary time-depth principle': implications for the reclassification of fear-circuitry-related traits in DSM-V and for studying resilience to war-zone-related posttraumatic stress disorder. *Prog Neuropsychopharmacol Biol Psychiatry* 2006; **30**:827–53.
- 6 Bracha HS, Williams AE, Haynes SN et al. The STRS (shortness of breath, tremulousness, racing heart, and sweating): a brief checklist for acute distress with panic-like autonomic indicators; development and factor structure. *Ann Gen Hosp Psychiatry* 2004; **3**:8.
- 7 Bracha HS, Garcia-Rill E, Mrak RE, Skinner RD. Post-mortem locus coeruleus neuron count in three American veterans with probable or possible war-related PTSD. *J Neuropsychiatry Clin Neurosci* 2005; **17**:503–9.
- 8 Stein DJ, Ipser JC, Balkom AJ. Pharmacotherapy for social phobia. *Cochrane Database Syst Rev* 2004; **4**:CD001206.
- 9 Bracha HS, Lenze SM. Why are we using surgery as a first line of treatment for an anxiety disorder? *Clin Auton Res* (in press).

Conflicts of interest: none declared.

Immotile acral melanocytes?

DOI: 10.1111/j.1365-2133.2006.07542.x

SIR, Eccrine poroma (EP) is a frequently encountered neoplasm which usually presents as a flesh-coloured nodule on the soles of the feet and palms of the hands. We have previously reported a pigmented facial EP.¹ We recently encountered two more patients with pigmented EP located on the abdomen and buttock, respectively. We are intrigued by the locations of the pigmented variant of this adnexal tumour. As this tumour appears predominantly on the distal extremities, it is surprising that very few reports have documented pigmented variants of EP occurring at these sites. In the biopsy specimens of pigmented EP, dendritic melanocytes (MCs) were usually seen among the tumour lobules. Several hypotheses have been proposed to explain the mechanism of MC colonization in the pigmented tumours. It is known that sweat duct primordia contain MCs during the 14th week of gestation that are lost later in embryonic development.² Therefore, it is possible that certain tumour-related factors may activate MCs in the sweat gland acrosyringium after fetal life. This hypothesis, however, fails to explain the scarcity of reports on acral pigmented EP. Another explanation is that the MCs in the epidermis surrounding the tumour might have migrated to the tumour nest during tumour proliferation via the secretion of tumour-related factors such as fibroblast growth factor or endothelin-1.^{1,3} This proposed mechanism may provide a better explanation for the observed clinical phenomenon. Interestingly, in clinical situations involv-